# **BIO-SUCCINIC ACID: AN ENVIRONMENT-FRIENDLY PLATFORM CHEMICAL**

# Sumit Sharma<sup>1</sup>, Saurabh Jyoti Sarma<sup>1\*</sup>, Satinder Kaur Brar<sup>2,3</sup>

<sup>1</sup>Department of Biotechnology, Bennett University, Greater Noida (UP), 201310, India <sup>2</sup>INRS-ETE, Université du Québec, 490, de la Couronne, Québec, Canada G1K 9A9 <sup>3</sup>Department of Civil Engineering, Lassonde School of Engineering, York University North York (Toronto), Canada, M3J 1P3.

Received on: 15.04.2020

**Revised on:** 09.05.2020

#### Abstract

Global research for biomass-based products is swelling gradually due to depletion of fossil-based raw material as well as their negative effects on the environment. The hazardous chemical processes used for production of valuable industrial chemicals are directly responsible for environmental pollution. Therefore, use of alternative renewable resources as feedstock to produce such chemicals is gaining popularity. Succinic acid (SA) is platform chemical which can be used to produce bulk industrial chemicals with huge global demand such as adipic acid, 1,4-butandiol, maleic anhydride. Non-hazardous biological fermentation process can be used to produce succinic acid, which can be further converted to these chemicals. The bio-based approach uses CO2 as supplement during the process and it replaces fossil-based raw materials; therefore, it is environment friendly. Mostly, pure sugars or sugars derived from crop residues or lignocellulosic materials are used to produce succinic acid. However, utilization of agricultural waste aromatic compounds to produce succinic acid has not been investigated in detail and not being implemented anywhere. Lignin waste management is a problem for the cellulosic bio-refineries and finding the way for its biological utilization is still in the stage of research and development. Unlike sugar metabolism, bioconversion of aromatic compounds is relatively complicated. Interestingly, bioconversion of agricultural waste aromatic compounds could be targeted towards production of succinic acid. The purpose of the present review is to highlight this possibility.

# 1. Introduction

Succinic Acid (SA) is a biochemical compound produced in almost all organisms, aerobic as well as anaerobic either as an intermediate or end product of their metabolic pathway. Knowing the positive impact of succinic acid on human health, Nobel Prize winner, Robert Koch popularized succinate by means of an industrial organic compound in food industries and generated interest intended for its high scale industrial production. Due to its impressive utilization to produce various industrial chemicals like 1, 4-butanediol, 2-pyrrolidinone, tetrahydrofuran, and various other conventional products like pigments, additives, detergents, pharmaceuticals intermediate, etc., succinic acid becomes a valuable chemical [1,2]. The production of SA was initiated by Agricola in 1546. Initially, it was extracted from Amber through dry distillation which releases acidic anhydride. At that time, the succinic acid used to recognize by odor only. Later, numerous chemical approaches of succinic acid extraction from Amber were developed such as extraction by solvents like petroleum ether, methanol, and ammonium carbonate. At the end of 18th-century, chemical-based industries started using fossils fuels (petroleum, coal and natural gases) or catalytic hydrogenation of maleic anhydride to produce succinic acid. However, due to the concern of their limited natural feedstock, and negative environmental impacts, biotechnology is now playing a vital role to develop environment friendly processes to produce succinic acid [3].

CONTACT \*Corresponding author: saurabh.sarma@bennett.edu.in

Color versions of one or more of the figures in this article can be found online at www.ste.org © 2020 Save the Environment



Accepted on: 15.05.2020

#### Keywords

Succinic acid, Platform chemical, Market potential, Lignin, Aromatic substrates. Bio-based succinic acid has various benefits over its chemically synthesized counterpart as it utilizes CO<sub>2</sub> during the process, renewable raw materials are used and produce fewer toxic byproducts. Moreover, succinic acid fermentation not only utilizes CO<sub>2</sub> but also produces three important products together i.e. ethanol, succinic acid, and diethyl succinate. Michigan biotechnology institute (USA) had isolated the first commercial micro-organism "Anaerobiospirillum succiniciproducens" as a natural succinic acid producer. Some other micro-organisms isolated for this purpose are Actinobacillus succinogenes, Mannheimia succiniciproducens and Basfia succiniciproducens [4]. Low product yield and low productivity are the major limitations of these natural producers. Metabolic engineering can resolve these problems by modifying the biosynthetic pathway as desired. Metabolic engineering can be applied to natural producers and or any other microorganism whose metabolic pathways are well studied such as Escherichia coli, Corynebacterium glutamicum, and Saccharomyces cerevisiae. However, E. coli is usually selected as the model organism for metabolic engineering to produce succinic acid as its genome, proteome, metabolome and culture conditions are well established. Thus, significant interest has been observed to develop metabolically engineered strain for succinic acid production and various technologies to transform succinic acid into industrially important derivatives [5, 6].

As the applications of succinic acid are very vast, so its demand for at commercial level is high. In the 1990s, total succinic acid was manufactured around 18,000 MT (metric tons) and estimated to increase with 27.4% CAGR to 768 million MT by the year 2025. Major markets spread over for succinic acid use are surfactants/detergents, anti-corrosion agents, flavoring agents or antimicrobial agents in foods, health industry including vitamins, antibiotics, etc. In 2009, *Bioamber* and *Biohub* are two major French corporations have begun pilot-plant for succinic acid production. They

used genetically engineered E. coli strains for this. Later in 2010, Myriant and MBI International started pilot plant production using Escherichia coli and Actinobacillus succinogenes, respectively in the United States. As the value of bio-based succinic acid production increases, at the end of 2010, German-based BASF and Dutch CSM firms also started bio-succinic acid production in Spain. BASF isolated their own novel rumen bacteria Basfia succiniproducens for succinic acid production. Later on, Japanese agencies (Mitsubishi and Ajinomoto) came up with scaling-up of succinic acid production by using the Corynebacterium strain [7, 8]. All of these companies use direct sugar sources like corn starch, sugarcane molasses even though there are other low-cost and abundant cellulosic biomass such as wood hydrolysate [9] also available for succinic acid production. Although these materials need to be pretreated to release carbohydrate molecules available, still they are less expensive than refined carbohydrates. Apart from cellulosic source, aromatic compounds source (lignin) have the potential for succinic acid production. This review signifies the susceptibility of the succinic acid as an important building block chemical.

### 2. Succinic acid: Production approaches

Succinic acid production methods through biological routes are well studied along with genetic level modifications. Biobased succinic acid production is non-hazardous and environmentally friendly while the petroleum-based method uses harmful chemicals for succinic acid and their derivatives development. Vaswani and Sudeep [10] defined both the chemical and biological processes for succinic acid synthesis. The chemical method starts with the conversion of maleic anhydride to succinic anhydride by hydrogenation with the aid of catalyst like Ni/Si alloy. After that, hydrolysis converts it into succinic acid. But hydrogenation causes toxicity and increases the cost of the whole process. However, the non-toxic biochemical approach uses glucose and  $CO_2$  as substrates to produce succinic acid (Figure 1).



Figure 1: Succinic acid production via a chemical and biological method.

There are various applications of succinic acid in the field of food and chemical industry. It is used to produce adipic acid, 1,4 butanediol, tetrahydrofuran, succinonitrile, succinimide, polyesters, resins, acidulants, etc. Figure 2 describes various applications of succinic acid. Moreover, it is utilized in detergent and surfactant formations, as chain extender, ionchelator, and foaming agent preparations. It is also used for making resins, inks, coating materials, plasticizers, liquors, plant growth stimulants, electrolyte bath additives, biodegradable plastics, anti-microbial agents, pH modifiers and flavoring agents.

As per the authors' knowledge, till now there is no bio-based



Figure 2: Applications of succinic acid as a platform chemical.

#### 3. Global market potential of succinic acid

The global market of succinic acid is expected to reach at around 701 million USD by 2021. The global demand for succinic acid is around 40,000 tons/year-45,000 tons/year. A

comparative analysis of the global market potential of succinic acid against other valuable industrial chemicals are presented in Figure 3. The major succinic acid manufacturers using petrochemical process are Kawasaki Kasei Chemicals, Linyi Lixing Chemical, Anhui Sunsing Chemicals, Mitsubishi Chemical, Nippon Shokubai, Gadiv Petrochemical, Astatech, R-Biopharm, Evonik, Thyssenkrupp [11]. On the other hand, major bio-based succinic acid manufacturers are BioAmber, Reverdia, and Succinity (Table 1). In India, Thirumalai Chemicals Ltd., A. B. Enterprises, Chemicals Chamber and others (shown in Table 2) are chemical-based manufacturers of succinic acid [12,13].



succinic acid producing company in India.

Figure 3: The global market potential of various industrial products.

### Table 1: Succinic acid key player in world.

Company Name	Location			
Shanghai Runkey Biotech Co., Ltd	China			
Linyi Lixing Chemical Co., Ltd.	China			
HeBei Oukai Biotechnology Co.,Ltd	China			
A & Z Food Additives Co., Ltd.	China			
ZSIKI KFT	Hungary			
BioAmber	United States			
Myriant	United States			
Succinity	Germany			
Reverdia	Italy			
Gadiv Petrochemical Industries Ltd.	Israel			
Nippon Shokubai	Japan			
Mitsubishi Chemical Holdings	Japan			
Kawasaki Kasei Chemicals Ltd.,	Japan			
BASF	Germany			
DSM	Netherlands			

### Table 2: Succinic acid key player in India.

Company Name	Location		
B Joshi Agrochem Pharma	Mumbai		
Thirumalai Chemicals Ltd.	Mumbai		
Alpha Chemika	Mumbai		
A. B. Enterprises	Mumbai		
Axiom Chemicals Pvt. Ltd.	Vadodra		
Meru Chem Pvt. Ltd.	Mumbai		
Antares Chem Private Limited	Mumbai		
Labdhi Chemicals	Mumbai		
Arnish Laborates Private Limited.	Mumbai		
Alpha Chem	Ambala		
Natural Organics India	Delhi		
Kapchem International	Mumbai		
Chemicals Chamber	Delhi		

### 31 Economic Value of Succinic Acid

The current bio-based production of compounds and polymers (excluding-fuel) is estimated to be around 50 million tons [14]. But still, chemical production from fossilfuels was significantly higher, which was accounted to be 330 MT in 2009 [15]. The market of petrol-based production of succinic acid was estimated to decline by 2019 because of its negative influence on the environment. Due to this, bio-based production processes are preferred for use in the pharmaceutical and cosmetics industries. Its non-toxicity and high purity help in taking care of the personal health of consumers [16]. Global market of succinic acid can be divided into different sectors such as, pharmaceuticals (15%), industrial (57.1%), food (13.07%) and others (13.92%).

Among various applications of succinic acid, the largest application is BDO (1, 4-butanediol) that boosted the international bio-succinic acid marketplace by about 17% in the year 2013. BDO is used as a solvent in many industries. It can be further converted to  $\gamma$ -butyrolactone (GBL) and tetrahydrofuran (THF). So, OMR industry analyst and Grand view research estimated the demand for succinic acid increases with CAGR of 35% till 2020.

On an industrial scale, butyrolactone is mainly manufactured from maleic anhydride. So, the replacement of maleic anhydride (MAN) by bio-succinic acid will take over this market, which is less expensive and environment-friendly. Polyurethane, a thermosetting polymer is another example that shows increases in the market possibility of bio-succinic acid. A new polyurethane class Non-isocyanate based polythenes (NIPUs) has recently been developed, which will also increase the demand for bio-succinic acid [17]. About 25,000 to 30,000 MT (metric tons) succinic acid produces annually and expected to rise 10% every year. The addressable market of succinic acid will replace mainly the maleic anhydride, adipic acid, phthalic anhydride market. Because it can be used as a substitute for these chemicals. The succinic acid price in the market relies upon its purity. productivity, yield, recovery process, and raw material cost.

The succinic acid produced from the organic process is usually sold at a price of USD 5.9 to 9 per kg.

32 Expansion of bio-based succinic acid productions There are various companies that are using bio-based production routes succinic commercial production of acid. Major companies producing bio-succinic acid are BASF, BioAmber, Mitsubishi, Myriant, DSM, Mitsui & Co, Succinity, Roquette Freres S.A, Purac, and Reverdia. Most of them are joint ventures (JV) except Myriant. The main biobased succinic acid production market was covered by BioAmber, Succinity, and Reverdia with the production capacity of 350 MT, 500 MT, and 300 MT respectively [18]. BioAmber, Reverdia, and Succinity were able to produce 3,800 MT in the year 2011. Further, the Myriant corporation also enhanced the production volume with a dramatic increase in their production units. The world's first largest bio-based succinic acid industry BioAmber is located in France and now able to produce succinic acid nearly 3,000 MT/year. All the bio-based succinic acid-producing companies are using sugars extracted from different crops or biomasses. Table 3 shows the foremost bio-based succinic acid-producing player in the succinic acid market along with the details of major production capacity, kind of substrates, and class of microbes used in production [19].

Company name	Location	Substrate	Microbe	Commercial Operation	Production (tons/year)
Bioamber (in JV with Mitsui)	Sarnia, Ontario, Canada	Corn, wheat, cassava, rice, sugarcane, sugar beets	Bacteria	August 2015	30,000
Myriant	Lake Providence, LA, USA	Sorghum grain, Ligno cellulosic sugar from corn stover	Bacteria	June 2013	14,000
Reverdia (Roquette- DSM JV)	Cassano Spinola, Italy	starch from corn	Yeast	December 2012	10,000
Succinity (BASF-Corbion Purac JV)	Succinity BASF-Corbion Purac JV) Barcelona, Spain sucrose, glucose		Bacteria	March 2014	10,000

Table 3: Bio-Based succinic acid-producing companies

# 4. Substrate and micro-organisms for succinic acid: Natural and Engineered

Corn waste, sugarcane molasses, wheat grains, whey, lignocellulosic biomass or pure sugars are commonly used substrates for fermentation. Glucose is the main carbon source used to study the metabolic rate of microbes. Natural succinic acid-producing microbes such as *Actinobacillus carcinogens, Anaerobicspirullum succiniproducens*, Mannheimia succinicproducens are restricted to glucose only, and cannot utilize a variety of substrates [20]. Genetically engineered micro-organisms such as *Corynebacterium glutamicum*, *E. coli*, and a new specific isolate *Basfia succinoproducens* were also investigated for their role in succinic acid production [21]. *Basfia succinoproducens* isolated from the rumen gut is a natural succinic acid-producing microbe. Different kinds of bacteria, natural and genetically engineered, were being used for getting a higher titer of succinic acid. Lignocellulosic sugar and direct pure sugar derivatives are the mostly used substrates to produce succinic acid. Some researchers suggested that corn cob, cane molasses, starch, and lignocellulosic biomass are the substrates useful for succinic acid production [22-41].

A comparative analysis of succinic acid yield and titer on different carbon sources and hydrolysates is depicted in Figures 4 and 5.



Figure 4: Comparative study of the bacterial fermentation of different sugar derivatives and other carbon sources for succinic acid production.



Figure 5: Comparative study of the bacterial fermentation of different biomass hydrolysates for succinic acid production.

# 5. Metabolic engineering strategies for succinic acid production

Glucose is the starting substrate for biosynthesis of succinic acid in microbial cells. Initially, glycolysis occurs till phosphoenolpyruvate (PEP) and pyruvate formation. When  $CO_2$  enters the system under anaerobic condition, it activates enzymes like phosphoenolpyruvate carboxylase (ppc), phosphoenolpyruvate carboxylase kinase (ppck), and pyruvate carboxylase (pyc). Their carboxylases catalyze PEP and pyruvate to produces oxaloacetate. At this stage, the citric acid cycle appears as a reverse second half citric acid cycle to produce succinic acid (Figure 6). Other byproducts such as, acetate, ethanol, lactate formate also start producing from pyruvate. Apart from that, keto-glutarate and malate form amino acid such as aspartate and glutamate. Figures 6 (A and B) described all probable products formed along with succinic acid-producing under anaerobic conditions. The theoretical yield measured for the succinic acid from glucose is 0.66g/g. One mole of CO<sub>2</sub> is used per mole of succinic acid obtained.



Figure 6: (A) Major metabolic routes to succinic acid production, (B) Bio-synthetic pathway of the succinic acid from glucose: Green branch of the pathway is representing succinic acid and white is for the by-product.

#### 6. Potential to use aromatic substrates

Mostly, sugar derivatives were used to produce succinic acid, but aromatic compounds and phenol derivatives have not been studied to produce succinic acid. However, the phenolic derivatives were studied for other industrial chemicals such as adipic acid, muconic acid, but not for succinic acid. The phenol degradation potential by different micro-organisms is shown in Figure 7 [42-50]. A higher degradation rate for aromatics like catechol, phenol, benzoic acid, p-coumarate, 4-hydroxy benzoate was achieved by the implementation of a genetic engineering approach. Researchers have studied the degradation of the aromatic into products like adipic acid, pyruvate, lactate, muconic acid by Pseudomonas putida. Vanillin, pyridine and di-carboxylic acid were produced from *Rhodococcus josti* [51-57]. This has been done by deleting gene of ldhA, pflB, pta and ackA and overexpression of pyc in E. coli and in other naturally succinic acid-producing microorganisms. Table 4 clearly differentiates the strategies used for the metabolic change in micro-organism for utilizing

diverse categories of phenolics. For muconic acid production, *PcaHG*, *CatBC* and *vdh* genes suppression and overexpression of the *AroY*, *AsbF*, *CatA* genes were accomplished in *P. putida*. So, there are possibilities for enhancing the succinic acid production from phenol derivatives by appropriate metabolic engineering strategies.



Figure 7: Comparative study of phenol degradation (mg/l) in bacteria for succinic acid production.

Sr. No.	Substrate	Micro-organism	Bio- Product	Gene deletion	Gene addition	Approach	Product/ yield	Refs
1.	Catechol, phenol, and benzoic acid using glucose as additional	Pseudomonas putida Kt2440	Muconic acid adipic acid, Nylon	catB	benAB CD, catA	Homologous recombination and plasmid insertion	13 g/L	[56]
2.	Catechol, phenol, and benzoic acid using glucose as additional	Corynebacterium glutamicum MA-2	Muconic acid	muconate cyclo isomerase (catB)	Catechol- 1,2-dioxy genase (catA)	Homologous recombination and plasmid insertion	85 g/L	[54]
3.	Benzoate, p-coumarate, phenol,4-hydroxy benzoate,	Pseudomonas putida Kt2440	Muconic acid, Adipic acid	Pca HG	AroY, Dmp KLMONP	Homologous recombination	13.5 g/L	[52]
4.	Benzoate, p-coumarate	Pseudomonas putida Kt2440	Pyruvate Lactate	aceEF	catBCA	Homologous recombination and plasmid insertion	41.17 g/g	[51]
5.	Wheat straw	Rhodococcus jostii RHA1	Vanillin	vdh	-	Homologous recombination	96 mg/L	[57]

 Table 4: Phenol derivatives used to produce various valued products.

# 7. Downstream processing techniques for succinic acid purification

The succinic acid comes in the market as a white crystalline powder form. When succinic acid is coming from the microbial broth, it should be clear and without any impurity especially, from hydrolysates which are having color, minerals and metal impurities in excess amount. Therefore, purification of the compound is very crucial for any kind of product that comes in the market. There are various methods established for succinic acid purification, some of them are patented processes developed by succinic acid-producing companies and some are research-based methods. Table 5 describes the purification process developed and currently used by bio-based succinic acid manufacturing companies.

Cost-effective downstream processes and clarification methods are always desirable to cut the purification cost. Three major categories of the purification are electrodialysis, salt precipitation and ammonia-based neutralization (Figure 8). These are mostly used due to their recyclability and low cost. Electrodialysis is done by ion-exchange membranes with electrolytic conductivity to transfer ions faster for separation of product. It can be reused several times and very effective in getting high yield [58]. In the salt precipitation process, calcium hydroxide is used for neutralization and precipitate formation. Then, the solid is dissolved by adding sulfuric acid and finally passed through ion-exchangers to get clarified product [59]. Third, ammonia neutralization is used to produce salts of succinic acid as di-ammonium succinate after the addition of liquor ammonia. It can be transformed into mono form through heating at 135°C with pressure 50 psi. Final crystals of acid are dissolved in water and purified again with ion exchangers [60]. In all the three processes, ion exchangers are used for extracting the final highly purified product. So that only the purified product will come without any other salt or metal ion impurities.

Company	Purification Process
BASF	Succinic acid is salted out first as a precipitate, then, passed through an ion-change column with a mild temperature near 50 to $60^{\circ}$ C.
BioAmber	Ammonia treatment to form di-ammonium succinate salt then heated to convert it into mono- ammonium succinate. Further dissolving in reverse osmosis water system.
Mitsubishi	Precipitation of succinic acid by reducing the overhead pressure below the atm. pressure.
Myriant	Succinic acid is purified from the vacuum dried concentrate of the broth, then acidic treatment separated it from ammonium sulfate.

Table 5: Succinic acid purification process developed by bio-succinic acid companies.



Figure 8: Various downstream processing methods for succinic acid.

#### 8 Conclusion

Succinic acid is well studied for its potential as a platform chemical, which has vast applications such as the formation of polymers. The process development for enhancement of yield and titer has been done by metabolic engineering. Due to the depletion of petroleum-based resources, alternate bio-based approaches are being explored. Easy sugar-based process for succinic acid production is developed and marketed; however, aromatic substrates are not evaluated yet for succinic acid production. Therefore, to compete with the increasing demand for polymers, this basic platform chemical needs to be produced in bulk quantity and preferably from biomass. Thus, succinic acid production from low-cost agro-industrial wastes has become a major research area.

#### Acknowledgement

Authors are thankful to Bennett University, Greater Noida for financial support to complete this work.

### References

- 1. Zeikus JG, Jain MK, Elankovan P. Biotechnology of succinic acid production and markets for derived industrial products. Applied Microbiology and Biotechnology. 1999 May 1;51(5):545-52.
- 2 Thakker C, Martínez I, San KY, Bennett GN. Succinate production in Escherichia coli. Biotechnology Journal. 2012 Feb;7(2):213-24.

- 3 Vávra N. The Chemistry of Amber-Facts, Findings and Opinions. Annalen des Naturhistorischen Museums in Wien. Serie A für Mineralogie und Petrographie, Geologie und Paläontologie, Anthropologie und Prähistorie. 2009 Jan 1:445-73.
- **4 Song H, Lee SY**. Production of succinic acid by bacterial fermentation. Enzyme and Microbial Technology. 2006 Jul 3;39(3):352-61.
- 5. Delhomme C, Weuster-Botz D, Kühn FE. Succinic acid from renewable resources as a C 4 building-block chemical—a review of the catalytic possibilities in aqueous media. Green Chemistry. 2009;11(1):13-26.
- **6** Cheng KK, Wang GY, Zeng J, Zhang JA. Improved succinate production by metabolic engineering. BioMed Research International. 2013;2013.
- 7. Nghiem NP, Kleff S, Schwegmann S. Succinic acid: technology development and commercialization. Fermentation 3 (2): 26.
- 8 Yuzbashev TV, Yuzbasheva EY, Laptev IA, Sobolevskaya TI, Vybornaya TV, Larina AS, Gvilava IT, Antonova SV, Sineoky SP. Is it possible to produce succinic acid at a low pH?. Bioengineered Bugs. 2011 Mar 1;2(2):115-9.
- 9. Lee PC, Lee S, Hong SH, Chang HN, Park SC. Biological conversion of wood hydrolysate to succinic

acid by *Anaerobiospirillum succiniciproducens*. Biotechnology Letters. 2003 Jan 1;25(2):111-4.

- **10.** Vaswani S. Bio-based succinic acid. SRI Consulting Review. 2010Dec.
- 11. https://www.openpr.com/news/1231350/2018-2023-Petro-based-Succinic-Acid-Market-By-Industry-Size-Share-Growth-Trends-Analysis-and-Forecast.html (accessed on 25/10/2019)
- 12 https://www.mordorintelligence.com/industryreports/ succinic- acid- market ( accessed on 02/11/2019).
- *13* https://www.tradeindia.com/manufacturers/succinic-acid.html (*accessed on 02/11/2019*).
- **14 Higson A.** NNFCC. Estimate of chemicals and polymers from renewable resources. 2010. NNFCC. Estimate of Fermentation Products. 2010.
- 15. Shen L, Haufe J, Patel MK. Product overview and market projection of emerging bio-based plastics PRO-BIP 2009. Report for European polysaccharide network of excellence (EPNOE) and European bioplastics. 2009 Jun;243.
- **16 Tan JP, Jahim JM, Harun S, Wu TY**. Overview of the Potential of Bio-Succinic Acid Production from Oil Palm Fronds. Journal of Physical Science. 2017 Jan 2;28.
- **17. Kathalewar MS, Joshi PB, Sabnis AS, Malshe VC**. Non-isocyanate polyurethanes: from chemistry to applications. RSC Advances. 2013;3(13):4110-29.
- 18 Weastra SR. WP. Determination of market potential for selected platform chemicals: itaconic acid, succinic acid, 2, 5-furandicarboxylic acid, Report for BioConSepT FP7 project, 2013.
- **19.** https://www.chemicals-technology.com/projects/ (accessed on25/11/2018)
- **20.** Song H, Lee SY. Production of succinic acid by bacterial fermentation. Enzyme and Microbial Technology. 2006 Jul 3;39(3):352-61.
- **21.** Li Q, Xing J. Microbial succinic acid production using different bacteria species. In: Microorganisms in Biorefineries 2015 (pp. 183-206). Springer, Berlin, Heidelberg.
- 22 Zhang W, Zhang T, Song M, Dai Z, Zhang S, Xin F, Dong W, Ma J, Jiang M. Metabolic engineering of Escherichia coli for high yield production of succinic acid driven by methanol. ACS Synthetic Biology. 2018 Oct 9;7(12):2803-11.
- 23. Duarte JC, Valença GP, Moran PJ, Rodrigues JA. Microbial production of Propionic and Succinic acid from Sorbitol using *Propionibacterium acidipropionici*. AMB Express. 2015 Dec 1;5(1):13.

- 24 Xu S, Hao N, Xu L, Liu Z, Yan M, Li Y, Ouyang P. Series fermentation production of ornithine and succinic acid from cane molasses by *Corynebacterium glutamicum*. Biochemical Engineering Journal. 2015 Jul 15;99:177-82.
- 25. Lu CD, Xiong YY, Zhao JF, Wang JH. Fermentation production of succinic acid from sucrose by engineered Escherichia coli JH208. In: Advanced Materials Research 2015 (Vol. 1088, pp. 503-506). Trans Tech Publications Ltd.
- 26 Carvalho M, Matos M, Roca C, Reis MA. Succinic acid production from glycerol by *Actinobacillus succinogenes* using dimethylsulfoxide as electron acceptor. New Biotechnology. 2014 Jan 25;31(1):133-9.
- **27.** Krishnakumar J. Biological production of succinic acid using a cull peach medium. 2013.
- 28 Leung CC, Cheung AS, Zhang AY, Lam KF, Lin CS. Utilisation of waste bread for fermentative succinic acid production. Biochemical Engineering Journal. 2012 Jun 15;65:10-5.
- 29. Okino S, Noburyu R, Suda M, Jojima T, Inui M, Yukawa H. An efficient succinic acid production process in a metabolically engineered *Corynebacterium glutamicum* strain. Applied Microbiology and Biotechnology. 2008 Dec 1;81(3):459-64.
- **30** Oh IJ, Lee HW, Park CH, Lee SY, Lee J. Succinic acid production by continuous fermentation process using *Mannheimia succiniciproducens* LPK7. Journal of Microbiology and Biotechnology 2008 May 1;18(5):908-12.
- **31.** Nghiem NP, Davison BH, Suttle BE, Richardson GR. Production of succinic acid by *Anaerobiospirillum* succiniciproducens. Applied Biochemistry and Biotechnology. 1997 Mar 1;63(1):565.
- Zhao Y, Cao W, Wang Z, Zhang B, Chen K, Ouyang P. Enhanced succinic acid production from corncob hydrolysate by microbial electrolysis cells. Bioresource Technology. 2016 Feb 1;202:152-7.
- 33 Salvachúa D, Smith H, John PC, Mohagheghi A, Peterson DJ, Black BA, Dowe N, Beckham GT. Succinic acid production from lignocellulosic hydrolysate by *Basfia succiniciproducens*. Bioresource Technology. 2016 Aug 1;214:558-66.
- 34 Shi X, Chen Y, Ren H, Liu D, Zhao T, Zhao N, Ying H. Economically enhanced succinic acid fermentation f rom cassava bagasse hydrolysate using *Corynebacterium glutamicum* immobilized in porous polyurethane filler. Bioresource Technology. 2014 Dec 1;174:190-7.

- 35. Liu R, Liang L, Li F, Wu M, Chen K, Ma J, Jiang M, Wei P, Ouyang P. Efficient succinic acid production from lignocellulosic biomass by simultaneous utilization of glucose and xylose in engineered *Escherichia coli*. Bioresource Technology. 2013 Dec 1;149:84-91.
- **36** Ke-Quan C, Han Z, Ye-Lian M, Min J, Jie-Yu C. Enhanced succinic acid production from sake lees hydrolysate by dilute sulfuric acid pretreatment and biotin supplementation. Journal of Sustainable Bioenergy Systems. 2012 Jun 18;2012.
- Borges ER, Pereira N. Succinic acid production from sugarcane bagasse hemicellulose hydrolysate by *Actinobacillus succinogenes*. Journal of Industrial Microbiology & Biotechnology. 2011 Aug 1;38(8):1001-11.
- 38 Chen KQ, Li J, Ma JF, Jiang M, Wei P, Liu ZM, Ying HJ. Succinic acid production by *Actinobacillus succinogenes* using hydrolysates of spent yeast cells and corn fiber. Bioresource Technology. 2011 Jan 1;102(2):1704-8.
- Li Q, Yang M, Wang D, Li W, Wu Y, Zhang Y, Xing J, Su Z. Efficient conversion of crop stalk wastes into succinic acid production by *Actinobacillus* succinogenes. Bioresource Technology. 2010 May 1;101(9):3292-4.
- **40** Zheng P, Dong JJ, Sun ZH, Ni Y, Fang L. Fermentative production of succinic acid from straw hydrolysate by *Actinobacillus succinogenes*. Bioresource Technology. 2009 Apr 1;100(8):2425-9.
- **41.** Kim DY, Yim SC, Lee PC, Lee WG, Lee SY, Chang HN. Batch and continuous fermentation of succinic acid from wood hydrolysate by *Mannheimia succiniciproducens* MBEL55E. Enzyme and Microbial Technology. 2004 Dec 1;35(6-7):648-53.
- 42 Alon N, Ron A, Eyal Z, J Stefan R, Charles L G. Phenol biodegradation by *Corynebacterium glutamicum* encapsulated in electrospun fibers. Journal of Environmental Protection. 2012 Feb 24;2012.
- **43 Pazarlioğlu NK, Telefoncu A**. Biodegradation of phenol by *Pseudomonas putida* immobilized on activated pumice particles. Process Biochemistry. 2005 Apr 1;40(5):1807-14.
- 44 Hinteregger C, Leitner R, Loidl M, Ferschl A, Streichsbier F. Degradation of phenol and phenolic compounds by *Pseudomonas putida* EKII. Applied Microbiology and Biotechnology. 1992 May 1;37(2):252-9.
- **45.** Mahiudddin M, Fakhruddin AN. Degradation of phenol via meta cleavage pathway by *Pseudomonas*

*fluorescens* PU1. ISRN Microbiology. 2012 Jan 23;2012.

- **46** Hakim LN, Rochmadi, Sutijan . Phenol biodegradation by immobilized *Pseudomonas putida* FNCC-0071 cells in alginate beads. In: AIP Conference Proceedings 2017 Jun 15 (Vol. 1855, No. 1, p. 050004). AIP Publishing LLC.
- **47.** Shumkova ES, Solyanikova IP, Plotnikova EG, Golovleva LA. Phenol degradation by *Rhodococcus opacus* strain 1G. Applied Biochemistry and Microbiology. 2009 Jan 1;45(1):43-9.
- **48** Westerberg K, Elväng AM, Stackebrandt E, Jansson JK. *Arthrobacter chlorophenolicus* sp. nov., a new species capable of degrading high concentrations of 4-chlorophenol. International Journal of Systematic and Evolutionary Microbiology. 2000 Nov 1;50(6):2083-92.
- 49. Sahoo NK, Pakshirajan K, Ghosh PK, Ghosh A. Biodegradation of 4-chlorophenol by *Arthrobacter chlorophenolicus* A6: effect of culture conditions and degradation kinetics. Biodegradation. 2011 Apr 1;22(2):275-86.
- **50 Hasan SA, Jabeen S**. Degradation kinetics and pathway of phenol by *Pseudomonas* and *Bacillus* species. Biotechnology & Biotechnological Equipment. 2015 Jan 2;29(1):45-53.
- **51. Johnson CW, Beckham GT**. Aromatic catabolic pathway selection for optimal production of pyruvate and lactate from lignin. Metabolic engineering. 2015 Mar 1;28:240-7.
- 52 Vardon DR, Franden MA, Johnson CW, Karp EM, Guarnieri MT, Linger JG, Salm MJ, Strathmann TJ, Beckham GT. Adipic acid production from lignin. Energy & Environmental Science. 2015;8(2):617-28.
- 53 Johnson CW, Salvachúa D, Khanna P, Smith H, Peterson DJ, Beckham GT. Enhancing muconic acid production from glucose and lignin-derived aromatic compounds via increased protocatechuate decarboxylase activity. Metabolic Engineering Communications. 2016 Dec 1;3:111-9.
- 54 Becker J, Kuhl M, Kohlstedt M, Starck S, Wittmann C. Metabolic engineering of *Corynebacterium glutamicum* for the production of cis, cis-muconic acid from lignin. Microbial Cell Factories. 2018 Dec;17(1):115.
- 55. Mycroft Z, Gomis M, Mines P, Law P, Bugg TD. Biocatalytic conversion of lignin to aromatic dicarboxylic acids in *Rhodococcus jostii* RHA1 by rerouting aromatic degradation pathways. Green Chemistry. 2015;17(11):4974-9.

- 56 Kohlstedt M, Starck S, Barton N, Stolzenberger J, Selzer M, Mehlmann K, Schneider R, Pleissner D, Rinkel J, Dickschat JS, Venus J. From lignin to nylon: cascaded chemical and biochemical conversion using metabolically engineered *Pseudomonas putida*. Metabolic Engineering. 2018 May 1;47:279-93.
- Sainsbury PD, Hardiman EM, Ahmad M, Otani H, Seghezzi N, Eltis LD, Bugg TD. Breaking down lignin to high-value chemicals: the conversion of lignocellulose to vanillin in a gene deletion mutant of *Rhodococcus jostii* RHA1. ACS Chemical Biology. 2013 Oct18;8(10):2151-6.
- Hongo M, Nomura Y, Iwahara M. Novel method of lactic acid production by electrodialysis fermentation. Applied and Environmental Microbiology. 1986 Aug 1;52(2):314-9.
- **59.** Datta R, Glassner DA, Jain MK, Roy JR. Fermentation and purification process for succinic acid. United States patent US 5,168,055. 1992 Dec 1.
- **60.** Huh YS, Jun YS, Hong YK, Song H, Lee SY, Hong WH. Effective purification of succinic acid from fermentation broth produced by *Mannheimia succiniciproducens*. Process Biochemistry. 2006 Jun 1;41(6):1461-5.