



## THE ADVERSE EFFECTS OF ENVIRONMENTAL NOISE EXPOSURE ON HEALTH

Manish Shukla<sup>1,2</sup> and Neeru Kapoor<sup>1,\*</sup>

<sup>1</sup>Defence Institute of Physiology and Allied Sciences, Delhi -110 054

<sup>2</sup>Department of Neurosurgery, Penn State College of Medicine, Milton S. Hershey Medical Centre, Hershey Pennsylvania USA;

Received on: 17.01.2022

Revised on: 23.01.2022

Accepted on: 05.02.2022

### Abstract

Noise has always been very common environmental problem for mankind. Noise leading to hearing loss is one of the most common neurological disorders. In defence scenario, armed forces are exposed to the following types of noise: tanks, aircrafts, bombardment, missile launching, ammunition trials etc. Noise incites both direct (hearing loss, tinnitus) as well as indirect effects (anxiety, depression, annoyance, etc.). Environmental noise has multiple effects on human health. Some of these effects such as raised blood pressure; hearing loss and cognitive impairments may have implications for adult health as well.

In this review, we consigned that noise affects memory storage, cognitive decision making, executive functions that are the foremost requirements for human being. Psychological behaviour like argumentativeness, mood fluctuations, incapability to face stressful situations hamper the performance and lead to a poor quality of life. Noise exposures disturb the balance of awake and sleep modes of brain activity leading to neurological change

### Keywords

Noise; Cognitive; Oxidative Stress; Hearing Loss.

### INTRODUCTION

Noise is defined as unwanted sound or a combination of sounds that has adverse effects on health. The effects can manifest in the form of physiologic damage or psychological harm through a variety of mechanisms (Bahng and Lee, 2020). WHO reported that around one million healthy life years are lost every year due to environmental noise. Noise Induced Hearing Loss (NIHL) is thought to be one of the major causes of preventable hearing loss. The psychological effects of noise are usually not well characterized and often ignored. However, their effect can be equally devastating and may include hypertension, tachycardia, elevated cortisol levels and increased physiologic stress. Collectively, these effects can have severe adverse consequences on daily living

and globally on economic production (Seidman and Standring, 2010). Currently the armed forces are facing severe disabilities secondary to noise. It has been shown that the top two disabilities now facing the American Military are hearing loss and tinnitus (Bahng and Lee, 2020). It is reported that the physical and emotional effects of persistent noise lead to irritability, anger, nausea, headache, sleep disturbances and higher sense of helplessness, lack of control, tension, stress, unhappiness, anxiety and depression (Fooladi, 2012). In a self-reported study on the uncontrolled noise exposures to honking and shouting, the participants indicated their responses as being rage and frustration (Fooladi, 2012). Noise may cause cognitive impairment by a variety of mechanisms. Earlier studies have demonstrated that children

in noisy environments have decreased attention on task and have lower performance on cognitive assignments compared to children in quiet environment (Hygge et al., 2003); Shield et al, 2015). A more recent study by Ljung et al., found that Traffic noise significantly impaired reading ability and comprehension as well as basic mathematics performance in children (Ljung et al, 2009). These psychological and physiological non-auditory effects of noise result in detrimental health consequence and a decreased quality of life (Seidman and Standring, 2010).

### **i. NOISE AND HEALTH IMPACT**

Increased arousal, learned helplessness, frustration, annoyance and consequences of sleep disturbance on performance are reported to contribute to cognitive changes (Stansfeld et al., 2000; (Evans, 2006). Noise exposure may slow rehearsal in memory, influence process of selectivity in memory and choice of strategies for carrying out a task (Smith et al, 1997). It may reduce helping behaviour, increase aggression and reduce the processing of social cues during task performance (Jones et al., 1981). Other physiological effects of noise exposure include symptoms such as nausea, headache, argumentativeness and change in mood and anxiety (Stansfeld et al., 2000). Noise induced disruption was also found for non- auditory tasks like serial recall of visually presented lists and reading (Klatte et al., 2013). Studies undertaken in rats by Rabat have shown that noise exposure was correlated to an incapability to face stressful situations (Rabat, 2007). Noise annoyance can result from noise interfering with daily activities, feelings, thoughts, sleep, or rest and might be accompanied by negative responses such as anger, displeasure, and exhaustion and by stress related symptoms (Tiesler et al., 2013).

An efficient evaluation of noise effect should include an analysis of its frequency spectrum (Mahendra Prashanth and Venugopalachar, 2011). (Shukla, 2012) showed that traffic noise produces both a high frequency component around 1 kHz and a low frequency component around 63 Hz, responsible for the health relevant hazards. Some of the problems of high frequency noise include hearing impairment, hypertension, high blood pressure, speech interference, annoyance and disturbance to daily activities whereas low frequency noise could lead to annoyance, sleep deprivation, physiological disorders, etc. (Davies and Kamp, 2012; Shukla, 2012). In another study, Prashanth et al. (2008) reported that repeated noise exposures in the low (> 22- 500 Hz) and mid frequency (>500 Hz to 2 kHz) octave bands lead to ear vibration, chronic fatigue, headache, awakening from sleep, neck pain, backache, eye ball pressure, and ear pulsation.

Animal Studies in Wistar rat show that these extra-auditory effects of noise exposure were due to modified levels of oxidative markers in two areas of the brain, mainly the hippocampus and cerebellum (Shukla et al., 2019; Uran et al., 2010). Significant decrease in the volume of granule layer and cells in cerebellum, decrease in somal volume of Purkinje cells and increased plasma corticosterone concentration was

found in noise exposed animals (Hosseini-Sharifabad and Sabahi, 2014).

### **ii. NOISE AND OXIDATIVE STRESS**

Certain environmental challenges can increase the production of reactive oxygen species (ROS) in different structures, which may override the cellular antioxidant defences and can lead to oxidative stress (Vicente et al., 2004). Interestingly, the abundance of polyunsaturated fatty acids and the low-level of defensive mechanisms, together with the high oxygen consumption, make the brain more susceptible to oxidative damage than other organs. In particular, after experimental noise exposure in laboratory animals, superoxide anion radicals emerge in the stria vascularis (Yamane et al., 1995), hydroxyl radicals significantly increase in the cochlea (Henderson et al., 2008), hydrogen peroxide-induced cell damage to the inner ear occurs in vitro (Dehne et al., 2000), glutathione increases in the lateral wall (Yamasoba et al., 1998) and glutathione peroxidase and malondialdehyde activities increase progressively with noise intensity in hair cells (Yamashita et al., 2004). Although experimental data show that loud noise can increase ROS in the auditory pathway (Yamashita et al., 2004; Pouyatos et al., 2005; Le Prell et al., 2007), reports concerning the influence of noise stress on oxidative status in extra-auditory CNS structures are scarce (Manikandan et al., 2006; Rabat et al., 2006; Turner et al., 2005).

In addition, experimental data in developing animals are lacking. In consequence, since the tonotopic organization of the auditory cortex is much more susceptible to perturbations of acoustic inputs in infancy than in older animals, as suggested by Wang studies (2004), the concept of a 'critical' or sensitive period of development after which plasticity is more restricted could be tested by exposing immature animals (Wang, 2004). The adverse effects of environmental noise on human mental health have been reported by several authors. In particular, it has been suggested that loud noise can induce a variety of symptoms, including changes in anxiety, emotional stress, increase in social conflicts, as well as general psychiatric disorders (Rabat et al., 2006). Similarly, animal experiments have demonstrated that acute and chronic noise exposures can also induce temporary or permanent changes related to the central nervous system (CNS) (Goble et al., 2009; Manikandan et al., 2006).

### **iii. NOISE AND HEARING LOSS**

Noise-induced hearing loss (NIHL) acquired in leisure or occupational settings is a common cause of hearing impairment in industrialized countries, with a prevalence second only to age-related hearing loss (ARHL) (Stanbury 2008). Hearing loss associated with mild acoustic overexposure is reversible, and hearing recovers within 2–3 weeks [Dengerink 1985]. This temporary loss is known as a temporary threshold shift (TTS), and is probably due to reversible damage to the stereocilia of hair cells [Gao 1992] and/or swelling, followed by recovery, of cochlear nerve terminals [Pujol 1999]. Noise intensity and duration of exposure determine the level of noise damage to an organism.

High-intensity noise exposure damages inner hair cells (IHCs), primarily through two pathways: direct mechanical damage, in which noise can destroy the static cilia of hair cells, resulting in hair cell loss and damage to supporting cells and spiral ganglia (Shukla et al., 2019)


#### iv. NOISE AND COGNITIVE FUNCTION

The damaging effect of noise, however, is not limited in the auditory system, but extend to many other systems (Basner et al., 2014). Recent studies have warned of noise-related impairment of learning ability and cognitive performance (Cheng et al., 2011; Cui et al., 2009; Wright et al., 2006). Studies suggest that the evidence of the effects of noise on children's cognition has grown stronger over recent years, with over 20 studies showing detrimental effects of noise on children's memory and reading outcomes [Belojevic

2012]. children exposed to chronic aircraft or road traffic noise at school have poorer reading comprehension and memory than children who are not exposed [Haines 2001].

#### v. HYPOTHESIZED RELATIONSHIPS BETWEEN HEARING LOSS AND COGNITIVE DECLINE

Several possible relationships have been postulated (see Figure 1). Cognitive decline may reduce the cognitive resources that are available for auditory perception, manifesting as hearing loss. In contrast, Lin and colleagues (Lin et al., 2013) have suggested that hearing loss causes cognitive decline that is either permanent (the “sensory-deprivation”), or potentially remediable (the “information-degradation”). Another possibility is that a third factor causes both declines (the “common cause”).



**Figure 1:** A summary of the four hypotheses for the causal relationship between hearing loss and cognitive decline. (A) Cognitive load on perception hypothesis: cognitive decline leads to audiometric hearing loss; (B) Information degradation hypothesis: audiometric hearing leads to diminished (but reversible) cognitive performance; (C) Sensory deprivation hypothesis: audiometric hearing loss causes cognitive decline; (D): Common cause hypothesis: both audiometric hearing loss and cognitive decline are caused by a third, common factor.

Soldiers who were exposed to unnecessary noise levels including explosions and blast waves shown severe NIHL and tinnitus (Kapoor 2019), as well as cognitive discrepancies and memory impairment (Belanger et al., 2009). The mechanisms underlying cognitive function decline after noise exposure are not entirely clear. Noise exposure impair cognitive functions through different pathway. One is related to the oxidative reaction initiated by noise exposure. Increased oxidative stress has been reported in many studies as the cause of neuronal degeneration seen in many auditory nuclei as well as in the brain regions critical for cognitive functions (Shukla 2019, 2020). The other way is due to change of auditory input to the cognitive brain after

hearing loss induced by noise. This latter approach has not been considered intensively in the past, but the possibility has been supported by the connection between the auditory brain and cognitive brain (Kraus and Canlon, 2012) and established by the hippocampal deterioration and deteriorated spatial memory in C57BL/6J mice with age related hearing loss (Yu et al., 2011) and the destruction of hippocampal neurogenesis in the rat after noise-induced unilateral hearing loss (Kraus et al., 2010) (Shukla 2019; 2020).

In the auditory system, environmental noise induces cell death (Basta, et al., 2005) and a threshold shift (Syka and Rybalko, 2000), and causes abnormal neural coding

(Pienkowski and Eggermont 2012) in the auditory cortex (AC) and other cortical regions responsible for acoustic processing (Cheng 2016). In contrast, environmental noise also affects non-auditory brain regions such as the hippocampus, a major limbic region that receives direct or indirect neural input from the central auditory system (Kraus and Canlon, 2012). Accumulating evidence illustrates a significant effect of noise on hippocampal-related cognition (Manikandan, et al., 2006). A number of studies including our own have highlighted that exposure to moderate noise can affect the physiological structure and subsequently the function of both the AC and hippocampus (Cheng, et al., 2011). However, does the AC, the primary brain region to process acoustic information, have high susceptibility to environmental noise than the hippocampus in non-auditory system? It has not been determined. Previous studies have shown that as a stressor, noise may cause extensive oxidative stress along the lemniscal ascending pathway including the AC and the hippocampus (Manikandan, et al., 2006; Cheng, et al., 2011). A moderate level of noise was used to avoid direct physical damage. The central nervous system undergoes progressive structural and functional maturation during postnatal development, and noise exposure, as a stressor, may therefore affect brain function of mice to a greater extent at this stage than in adults (Chang and Merzenich, 2003).

#### vi. NOISE AND CARDIOVASCULAR RISK

Both short-term laboratory studies of human beings and long-term studies of animals have provided biological mechanisms and plausibility for the theory that long-term exposure to environmental noise affects the cardiovascular system and causes manifest diseases (including hypertension, ischaemic heart diseases, and stroke) (Basner 2014). Acute exposure to different kinds of noise is associated with arousals of the autonomic nervous system and endocrine system (Lusk 2004). Investigators have repeatedly noted that noise exposure increases systolic and diastolic blood pressure, changes heart rate, and causes the release of stress hormones (including catecholamines and glucocorticoids) Basner 2014. The general stress model is the rationale behind these reactions. Potential mechanisms are emotional stress reactions due to perceived discomfort (indirect pathway), and non-conscious physiological stress from interactions between the central auditory system and other regions of the CNS (direct pathway). The direct pathway might be the predominant mechanism in sleeping individuals, even at low noise levels.

Chronic exposure can cause an imbalance in an organism's homeostasis (allostatic load), which affects metabolism and the cardiovascular system, with increases in established cardiovascular disease risk factors such as blood pressure, blood lipid concentrations, blood viscosity, and blood glucose concentrations (Basner 2014; Babisch 2011). These changes increase the risk of hypertension, arteriosclerosis, and are related to severe events, such as myocardial infarction and stroke. Studies of occupational (Davies 2012) and environmental epidemiology have shown a higher

prevalence and incidence of cardiovascular diseases and mortality in highly noise-exposed groups (Sørensen 2012).

#### CONCLUSION

In summary, there is sufficient evidence for the effects of environmental noise in human being on hormonal changes, annoyance, well-being and cognitive effects such as reading comprehension, long-term memory and performance. The review finding that aircraft/environmental noise is related to hyperactivity symptoms requires some systematic investigation and undoubtedly the prolonged use of personal listening devices on hearing needs to be assessed in longitudinal studies, not least because of the public health implications of almost universal use in young people.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### REFERENCES

1. **Bahng, J., Lee, C.H.,** 2020. Topic Modeling for Analyzing Patients' Perceptions and Concerns of Hearing Loss on Social Q&A Sites: Incorporating Patients' Perspective. *Int. J. Environ. Res. Public Health* 17, 6209. <https://doi.org/10.3390/ijerph17176209>
2. **Basner, M., Babisch, W., Davis, A., Brink, M., Clark, C., Janssen, S., Stansfeld, S.,** 2014. Auditory and non-auditory effects of noise on health. *Lancet* (London, England) 383, 1325–32. [https://doi.org/10.1016/S0140-6736\(13\)61613-X](https://doi.org/10.1016/S0140-6736(13)61613-X).
3. **Cheng, L., Wang, S.-H., Chen, Q.-C., Liao, X.-M.,** 2011. Moderate noise induced cognition impairment of mice and its underlying mechanisms. *Physiol. Behav.* 104, 981–8. <https://doi.org/10.1016/j.physbeh.2011.06.018>.
4. **Chengzhi, C., Yan, T., Xuejun, J., Xiang, L., Youbin, Q., Baijie, T.,** 2011. Recovery of chronic noise exposure induced spatial learning and memory deficits in young male Sprague-Dawley rats. *J. Occup. Health* 53, 157–63. <https://doi.org/10.1539/joh.L10125>.
5. **Cui, B., Wu, M., She, X.,** 2009. Effects of chronic noise exposure on spatial learning and memory of rats in relation to neurotransmitters and NMDAR2B alteration in the hippocampus. *J. Occup. Health* 51, 152–8.
6. **Davies, H., Kamp, I.,** 2012. Noise and cardiovascular disease: A review of the literature 2008-2011. *Noise Heal.* 14, 287. <https://doi.org/10.4103/1463-1741.104895>.
7. **Evans, G.W.,** 2006. Child Development and the Physical Environment. *Annu. Rev. Psychol.* 57, 423–451. <https://doi.org/10.1146/annurev.psych.57.102904.190057>.
8. **Evans GW, Hygge S.** Noise and performance in children and adults. In: Luxon L, Prasher D, editors. *Noise and its effects*. London: Whurr Publishers; 2007.
9. **Fooladi, M.M.,** 2012. Involuntary and Persistent Environmental Noise Influences Health and Hearing in

- Beirut, Lebanon. *J. Environ. Public Health* 2012, 1–7. <https://doi.org/10.1155/2012/235618>.
10. **Goble, T.J., Møller, A.R., Thompson, L.T.,** 2009. Acute high-intensity sound exposure alters responses of place cells in hippocampus. *Hear. Res.* 253, 52–9. <https://doi.org/10.1016/j.heares.2009.03.002>.
  11. **Henderson, D., Hu, B., Bielefeld, E.,** 2008. Patterns and Mechanisms of Noise-Induced Cochlear Pathology. *Audit. Trauma, Prot. Repair* 195–217. [https://doi.org/10.1007/978-0-387-72561-1\\_7](https://doi.org/10.1007/978-0-387-72561-1_7).
  12. **Hosseini-Sharifabad, M., Sabahi, A.,** 2014. Stereological estimation of granule cell number and purkinje cell volume in the cerebellum of noise-exposed young rat. *Iran. J. Med. Sci.* 39, 387–90.
  13. **Haines MM, Stansfeld SA, Job RF,** et al. Chronic aircraft noise exposure, stress responses, mental health and cognitive performance in school children. *Psychol Med.* 2001;31:265–77.
  14. **Hygge, S., Boman, E., Enmarker, I.,** 2003. The effects of road traffic noise and meaningful irrelevant speech on different memory systems. *Scand. J. Psychol.* 44, 13–21. <https://doi.org/10.1111/1467-9450.00316>.
  15. **Jones, D.M., Chapman, A.J., Auburn, T.C.,** 1981. Noise in the environment: A social perspective. *J. Environ. Psychol.* 1, 43–59. [https://doi.org/10.1016/S0272-4944\(81\)80017-5](https://doi.org/10.1016/S0272-4944(81)80017-5).
  16. **Klatte, M., Bergström, K., Lachmann, T.,** 2013. Does noise affect learning? A short review on noise effects on cognitive performance in children. *Front. Psychol.* 4. <https://doi.org/10.3389/fpsyg.2013.00578>.
  17. **Kraus, K.S., Canlon, B.,** 2012. Neuronal connectivity and interactions between the auditory and limbic systems. Effects of noise and tinnitus. *Hear. Res.* 288, 34–46. <https://doi.org/10.1016/j.heares.2012.02.009>.
  18. **Kraus, K.S., Mitra, S., Jimenez, Z., Hinduja, S., Ding, D., Jiang, H., Gray, L., Lobarinas, E., Sun, W., Salvi, R.J.,** 2010. Noise trauma impairs neurogenesis in the rat hippocampus. *Neuroscience* 167, 1216–26. <https://doi.org/10.1016/j.neuroscience.2010.02.071>.
  19. **Lin, F.R., Yaffe, K., Xia, J., Xue, Q.-L., Harris, T.B., Purchase-Helzner, E., Satterfield, S., Ayonayon, H.N., Ferrucci, L., Simonsick, E.M.,** Health ABC Study Group, for the, 2013. Hearing Loss and Cognitive Decline in Older Adults. *JAMA Intern. Med.* 173, 293. <https://doi.org/10.1001/jamainternmed.2013.1868>.
  20. **Ljung, Robert, Patrik Sorqvist, and Staffan Hygge.** "Effects of road traffic noise and irrelevant speech on children's reading and mathematical performance." *Noise and Health* 11.45 (2009): 194.
  21. **Mahendra Prashanth, K., Venugopalachar, S.,** 2011. The possible influence of noise frequency components on the health of exposed industrial workers - A review. *Noise Heal.* 13, 16. <https://doi.org/10.4103/1463-1741.73996>.
  22. **Manikandan, S., Padma, M.K., Srikumar, R., Jeya Parthasarathy, N., Muthuvel, A., Sheela Devi, R.,** 2006. Effects of chronic noise stress on spatial memory of rats in relation to neuronal dendritic alteration and free radical-imbalance in hippocampus and medial prefrontal cortex. *Neurosci. Lett.* 399, 17–22. <https://doi.org/10.1016/j.neulet.2006.01.037>.
  23. **Prashanth, KV Mahendra, and V. Sridhar.** "The relationship between noise frequency components and physical, physiological and psychological effects of industrial workers." *Noise and Health* 10.40 (2008): 90.
  24. **Rabat, A., Bouyer, J., George, O., Lemoal, M., Mayo, W.,** 2006. Chronic exposure of rats to noise: Relationship between long-term memory deficits and slow wave sleep disturbances. *Behav. Brain Res.* 171, 303–312. <https://doi.org/10.1016/j.bbr.2006.04.007>.
  25. **Rabat, Arnaud.** "Extra-auditory effects of noise in laboratory animals: the relationship between noise and sleep." *Journal of the American association for laboratory animal science* 46.1 (2007): 35-41.
  26. **Seidman, M.D., Standing, R.,** 2010. Noise and Quality of Life. *Int. J. Environ. Res. Public Health* 7, 3730–3738. <https://doi.org/10.3390/ijerph7103730>.
  27. **Shield, Bridget,** et al. "A survey of acoustic conditions and noise levels in secondary school classrooms in England." *The Journal of the Acoustical Society of America* 137.1 (2015): 177-188.
  28. **Smith, Andrew,** et al. "Effects of caffeine and noise on mood, performance and cardiovascular functioning." *Human Psychopharmacology: Clinical and Experimental* 12.1 (1997): 27-33.
  29. **Shukla,** 2012. Characterization of traffic noise for a typical Indian road crossing. *Curr. Sci.* 103, 1193–1201. <https://doi.org/10.2307/24089232>.
  30. **Shukla, M., Roy, K., Kaur, C., Nayak, D., Mani, K.V., Shukla, S., Kapoor, N.,** 2019. Attenuation of adverse effects of noise induced hearing loss on adult neurogenesis and memory in rats by intervention with Adenosine A2A receptor agonist. *Brain Res. Bull.* 147, 47–57. <https://doi.org/10.1016/j.brainresbull.2019.02.006>.
  31. **Stansfeld, S., Haines, M., Brown, B.,** 2000. Noise and Health in the Urban Environment. *Rev. Environ. Health* 15. <https://doi.org/10.1515/REVEH.2000.15.1-2.43>.
  32. **Tiesler, C.M.T., Birk, M., Thiering, E., Kohlböck, G., Koletzko, S., Bauer, C.-P., Berdel, D., von Berg, A., Babisch, W., Heinrich, J.,** 2013. Exposure to road traffic noise and children's behavioural problems and sleep disturbance: Results from the GINIplus and LISAPLUS studies. *Environ. Res.* 123, 1–8. <https://doi.org/10.1016/j.envres.2013.01.009>.
  33. **Turner, J.G., Parrish, J.L., Hughes, L.F., Toth, L.A., Caspary, D.M.,** 2005. Hearing in laboratory animals: strain differences and nonauditory effects of noise. *Comp. Med.* 55, 12–23.

34. **Uran, S.L., Caceres, L.G., Guelman, L.R.,** 2010. Effects of loud noise on hippocampal and cerebellar-related behaviors. *Brain Res.* 1361, 102–114. <https://doi.org/10.1016/j.brainres.2010.09.022>.
35. **Vicente, É., Boer, M., Netto, C., Fochesatto, C., Dalmaz, C., Rodrigues Siqueira, I., Gonçalves, C.-A.,** 2004. Hippocampal antioxidant system in neonates from methylmercury-intoxicated rats. *Neurotoxicol. Teratol.* 26, 817–823. <https://doi.org/10.1016/j.ntt.2004.08.003>.
36. **Wang, X.,** 2004. The unexpected consequences of a noisy environment. *Trends Neurosci.* 27, 364–366. <https://doi.org/10.1016/j.tins.2004.04.012>.
37. **Wright, R.L., Lightner, E.N., Harman, J.S., Meijer, O.C., Conrad, C.D.,** 2006. Attenuating corticosterone levels on the day of memory assessment prevents chronic stress-induced impairments in spatial memory. *Eur. J. Neurosci.* 24, 595–605. <https://doi.org/10.1111/j.1460-9568.2006.04948.x>.
38. **Yamashita, D., Jiang, H.-Y., Schacht, J., Miller, J.M.,** 2004. Delayed production of free radicals following noise exposure. *Brain Res.* 1019, 201–209. <https://doi.org/10.1016/j.brainres.2004.05.104>.
39. **Dehne, N.,** et al. "In vitro effects of hydrogen peroxide on the cochlear neurosensory epithelium of the guinea pig." *Hearing research* 143.1-2 (2000): 162-170.
40. **Pouyatos, Benoît, Caroline A. Gearhart, and Laurence D. Fechter.** "Acrylonitrile potentiates hearing loss and cochlear damage induced by moderate noise exposure in rats." *Toxicology and applied pharmacology* 204.1 (2005): 46-56.
41. **Le Prell, Colleen G.,** et al. "Mechanisms of noise-induced hearing loss indicate multiple methods of prevention." *Hearing research* 226.1-2 (2007): 22-43.
42. **Yamasoba, Tatsuya,** et al. "Role of glutathione in protection against noise-induced hearing loss." *Brain research* 784.1-2 (1998): 82-90.
43. **Stanbury, Martha, Ann P. Rafferty, and Kenneth Rosenman.** "Prevalence of hearing loss and work-related noise-induced hearing loss in Michigan." *Journal of Occupational and Environmental Medicine* (2008): 72-79.
44. **Pujol, Rémy, and Jean-luc Puel.** "Excitotoxicity, synaptic repair, and functional recovery in the mammalian cochlea: a review of recent findings." *Annals of the New York Academy of Sciences* 884.1 (1999): 249-254.
45. **Dengerink, Harold,** et al. "The recovery of vascular changes following brief noise exposure." *Acta otolaryngologica* 100.1-2 (1985): 19-25.
46. **Gao, Wen-yuan,** et al. "A comparison of changes in the stereocilia between temporary and permanent hearing losses in acoustic trauma." *Hearing research* 62.1 (1992): 27-41.
47. **Belojevic, Goran,** et al. "Traffic noise and executive functioning in urban primary school children: The moderating role of gender." *Journal of environmental psychology* 32.4 (2012): 337-341.
48. **Haines, Mary M.,** et al. "Chronic aircraft noise exposure, stress responses, mental health and cognitive performance in school children." *Psychological medicine* 31.2 (2001): 265-277.
49. **Kapoor, Neeru, K. V. Mani, and Manish Shukla.** "Distortion product oto-acoustic emission: A superior tool for hearing assessment than pure tone audiometry." *Noise & Health* 21.101 (2019): 164.
50. **Belanger, Heather G.,** et al. "Cognitive sequelae of blast-related versus other mechanisms of brain trauma." *Journal of the International Neuropsychological Society* 15.1 (2009): 1-8.
51. **Shukla, Manish,** et al. "Moderate Noise associated oxidative stress with concomitant memory impairment, neuro-inflammation and Neurodegeneration." *Brain, Behavior, & Immunity-Health.* 5 (2020): 100089.
52. **Yu, Ya-Feng,** et al. "The relationship between age-related hearing loss and synaptic changes in the hippocampus of C57BL/6J mice." *Experimental gerontology* 46.9 (2011): 716-722.
53. **Cheng, Liang,** et al. "The hippocampus may be more susceptible to environmental noise than the auditory cortex." *Hearing research* 333 (2016): 93-97.
54. **Pienkowski, Martin, and Jos J. Eggermont.** "Reversible long-term changes in auditory processing in mature auditory cortex in the absence of hearing loss induced by passive, moderate-level sound exposure." *Ear and hearing* 33.3 (2012): 305-314.
55. **Syka, Josef, and Natalia Rybalko.** "Threshold shifts and enhancement of cortical evoked responses after noise exposure in rats." *Hearing research* 139.1-2 (2000): 59-68.
56. **Chang, Edward F., and Michael M. Merzenich.** "Environmental noise retards auditory cortical development." *science* 300.5618 (2003): 498-502.
57. **Basta, Dietmar, Barbara Tzschentke, and Arne Ernst.** "Noise-induced cell death in the mouse medial geniculate body and primary auditory cortex." *Neuroscience letters* 381.1-2 (2005): 199-204.
58. **Lusk, Sally L.,** et al. "Acute effects of noise on blood pressure and heart rate." *Archives of Environmental Health: An International Journal* 59.8 (2004): 392-399.
59. **Babisch, Wolfgang.** "Cardiovascular effects of noise." *Noise and health* 13.52 (2011): 201.
60. **Davies, Hugh, and Irene Van Kamp.** "Noise and cardiovascular disease: A review of the literature 2008-2011." *Noise and Health* 14.61 (2012): 287.
61. **Sørensen, Mette,** et al. "Road traffic noise and incident myocardial infarction: a prospective cohort study." *PLoS one* 7.6 (2012): e39283.